



JOINT PRESERVATION INSTITUTE

Patellar Tendon Repair Post-Operative Protocol

Phase I – Maximum Protection (Weeks 0 to 6):

Weeks 0 to 2:

- Brace locked in full extension for 6 weeks
- Non-weight bearing for 2 weeks with brace on.
- Ice and modalities to reduce pain and inflammation
- Aggressive patella mobility drills
- Range of motion – 0° to 30° knee flexion
- Begin submaximal quadriceps setting

Weeks 2 to 4:

- 50 lbs Partial Weight bearing with brace on.
- Continue with inflammation control
- Continue with aggressive patella mobility
- Range of motion – 0° to 60°
- Continue with submaximal quadriceps setting

Weeks 4 to 6:

- Gradual progression of weight bearing to 100 lbs.
- Continue with ice and aggressive patella mobility
- Range of motion – 0° to 90° (by week 6)
- Increase intensity with quadriceps setting

Phase II – Progressive Range of Motion and Early Strengthening (Weeks 6 to 12):

Weeks 6 to 8:

- Full weight bearing
- Open brace to 45°- 60° of flexion week 5, 90° at week 6
- Continue with swelling control and patella mobility

- Gradually progress to full range of motion
- Begin multi-plane straight leg raising and closed kinetic chain strengthening program focusing on quality VMO function.
- Initiate open kinetic chain limited to 50 lbs.
- Normalize gait pattern
- Begin stationary bike
- Initiate pool program

Weeks 8 to 10:

- Wean out of brace
- Continue with patella mobility drills
- Normalize gait pattern
- Restore full ROM
- Progress open and closed kinetic chain program from bilateral to unilateral
- Increase intensity on stationary bike
- Begin treadmill walking program forward and backward
- Begin elliptical trainer

Weeks 10 to 12:

- Full ROM
- Terminal quadriceps stretching
- Advance unilateral open and closed kinetic chain strengthening

Phase III – Progressive Strengthening (Weeks 12 to 16):

Weeks 12 to 16:

- Advance open and closed kinetic chain strengthening
- Increase intensity on bike, treadmill, and elliptical trainer
- Increase difficulty and intensity on proprioception drills
- Begin gym strengthening: leg press, hamstring curls, ab/adduction; avoid lunges and knee extensions
- Begin multi-directional functional cord program

Phase IV – Advanced Strengthening and Functional Drills (Weeks 16 to 20):

Weeks 16 to 20:

- May begin leg extensions; 30° to 0° progressing to full ROM as patellofemoral arthrokinematics normalize
- Begin pool running program advancing to land as tolerated

Phase V – Plyometric Drills and Return to Sport Phase (Weeks 20 to 24):

Weeks 20 to 24:

- Advance gym strengthening
- Progress running/sprinting program
- Begin multi-directional field/court drills
- Begin bilateral progressing to unilateral plyometric drills
- Follow-up appointment with physician
- Sports test for return to competition